



Dr. Berg®

THE KNOWLEDGE DOC®

POSTMENOPAUSAL MUSCLE LOSS

RECOVERY GUIDE

Complete
3-part
action plan
—See inside





Dr. Berg®

THE KNOWLEDGE DOC®

The **get-it-back guide** to postmenopausal muscle loss

Use this guide as a go-to reference while you build long-term tissue strength and function. The goal isn't to hit the ground running. Doing too much too soon can increase your risk of injury and burnout.

A safer, more sustainable approach: small, consistent changes that compound into lasting improvements in strength, muscle health, and confidence.

First, build a nutritional foundation to support healthy tissue development. Then gradually add training that puts those nutrients to work.

This copy is for informational purposes only and is not intended to diagnose, treat, cure, or prevent any disease. Consult your physician before starting any new health regimen or supplement.



Your 3-Part Action Plan

Muscle loss tends to accelerate after menopause, in part due to hormone-driven anabolic resistance, in which muscle becomes less responsive to protein and exercise over time.

However, the responsiveness hasn't disappeared—your muscles just require a stronger physical and nutritional stimulus to trigger it.

This is why proper nutrition and targeted exercise are especially important for supporting muscle building and tissue repair after menopause.

The plan below focuses on the three primary steps that support muscle development and maintenance.

Part 1 — Protein

Muscle repair and growth depend on a steady supply of high-quality protein and leucine, and can be further supported by meal timing.

Step 1: Your daily protein target

While individual needs vary, women should generally aim for 0.54 to 0.72 grams of protein per pound of body weight daily.

For highly active or athletic women, [research](#) published in the *Journal of the International Society of Sports Nutrition* suggests that daily needs are a bit higher—roughly 0.64 to 1.0 gram of protein per pound of body weight.

Those looking to actively build or recover muscle may need to aim for the upper end of their target range.



Here is what a reasonable range looks like for the average woman:

Your weight	Daily protein target	Avg. per-meal target (2 meals)	Avg. per-meal target (3 meals)
120 lb (54 kg)	65–87 g/day	~33–44 g × 2 meals	~22–29 g × 3 meals
140 lb (64 kg)	76–102 g/day	~38–51 g × 2 meals	~25–34 g × 3 meals
160 lb (73 kg)	87–116 g/day	~44–58 g × 2 meals	~29–39 g × 3 meals
180 lb (82 kg)	98–131 g/day	~49–66 g × 2 meals	~33–44 g × 3 meals
200 lb (91 kg)	109–145 g/day	~55–73 g × 2 meals	~36–48 g × 3 meals

Here's what protein intake should look like for physically active women and female athletes:

Your weight	Daily protein target	Avg. per-meal target (2 meals)	Avg. per-meal target (3 meals)
120 lb (54 kg)	77–120 g/day	~39–60 g × 2 meals	~26–40 g × 3 meals
140 lb (64 kg)	90–140 g/day	~45–70 g × 2 meals	~30–47 g × 3 meals
160 lb (73 kg)	102–160 g/day	~51–80 g × 2 meals	~34–53 g × 3 meals
180 lb (82 kg)	115–180 g/day	~58–90 g × 2 meals	~38–60 g × 3 meals
200 lb (91 kg)	128–200 g/day	~64–100 g × 2 meals	~43–67 g × 3 meals



Step 2: Hit the leucine threshold

Total daily protein intake is only part of the picture. Each meal also needs to deliver enough leucine. Leucine is the amino acid that triggers the body's "build muscle" signal.

After menopause, the amount of leucine needed to trigger that signal rises significantly—up to twice the standard recommendation for younger adults.

To activate muscle protein synthesis after menopause, aim for roughly 3 to 4 grams of leucine per meal.

The following chart outlines some of the most concentrated sources of leucine.

Food	Serving	Leucine (approx)
Swiss cheese	1 cup	~3.9 g
Yellowtail fish	½ fillet	~3.5 g
Ham	1 cup	~3.1 g
Chicken drumstick meat	1 cup	~3.0 g
Whey protein isolate	Per ~35 g scoop	~3.0 g
Turkey	1 cup	~2.8 g
Lean beef (sirloin/round)	3 oz	~2.6 g
Cottage cheese	1 cup	~2.5 g
Canned tuna	3 oz	~2.0 g
Pork loin, cooked	3 oz	~2.0 g
Whole milk mozzarella	1 cup	~2.0 g
Salmon, cooked	3 oz	~1.8 g
Firm tofu	½ cup	~1.7 g
Pumpkin seeds	½ cup	~1.4 g
Pacific cod	1 fillet	~1.4 g
Parmesan/hard cheese	½ cup	~1.4 g
Whole milk yogurt, plain	12 oz	~1.2 g
Whole eggs	2 large	~1.1 g

Rule of thumb: According to [research](#) published in The American Journal of Clinical Nutrition, older adults should aim for at least 25 to 30 grams of quality protein per meal to stimulate muscle protein synthesis.



Step 3: Meal timing

Here are three meal-timing tips to help you support muscle growth and retention.

1. Avoid snacking

Efficient muscle building depends on blood leucine rising above a threshold of around 3 to 4 grams in a single meal. Grazing on small amounts of protein can keep you below that threshold, so the muscle-building signal may not fully activate.

2. Space out your meals

Continuous protein intake doesn't necessarily translate into consistent muscle growth. Once enough leucine triggers the anabolic response, your cells need about 3 to 4 hours to process nutrients, reset, and become sensitive to the next protein-rich meal.

As a result, protein eaten during this window may not trigger a strong building signal. Instead, it may be best to wait until after the resting period to enjoy your next protein-rich meal.

3. Choose the right fasting strategy

Many women use fasting to address metabolic changes such as slower metabolism, weight gain, and insulin resistance after menopause.

Fasting can support metabolic health, but longer windows—like one meal a day (OMAD)—can make it harder to meet protein needs and maximize muscle-building opportunities.

Strategic fasting while building muscle

To help protect lean mass and support muscle signaling while fasting, consider adopting one of the two protocols below.

Track A: Muscle maintenance

- **The target:** 2 to 3 quality, protein-anchored meals per day, spaced 3 to 4 hours apart, to meet your average protein target.
- **Acceptable fasting methods:** 12:12 through 18:6 time-restricted eating (TRE) patterns. These windows allow time for up to three threshold-crossing meals.



- **One meal a day (OMAD):** If preferred, only implement OMAD 2 to 3 days a week, not daily—and monitor how your strength and energy respond. On OMAD days, ensure the single meal is very high in protein.
- **Prolonged fasting (multiday fasts):** May not be ideal for women whose primary goal is maintaining or rebuilding muscle. These fasts may contribute to lean mass loss, so prioritize adequate protein intake and resistance training before and after fasting, and avoid stacking them back-to-back.

Track B: Rebuilding

- **The target:** Three protein-anchored meals per day, spaced 3 to 4 hours apart, aim for the higher end of your average protein requirement.
- **Acceptable fasting methods:** 12:12 up to 15:9 TRE or a standard lifestyle eating schedule. Avoid longer, aggressive fasting windows here.
- **OMAD:** Not recommended
- **Prolonged fasting:** Not recommended





Part 2 – Supporting nutrients

Beyond protein, several nutrients influence how well your muscles respond to physical training.

Below are four key nutrients for supporting normal muscle health and development. Potential sources are listed; use supplements only if appropriate for you.

Nutrient	Role in muscle development	Context/recommendation	Sources
Creatine	Helps supply energy during resistance training and supports strength and muscle adaptations. May play a role in supporting muscle recovery.	Often used in monohydrate form	Beef, pork, herring, salmon
Vitamin D	Supports normal muscle function. Low levels have been linked to greater muscle loss.	Get your levels tested and aim to maintain a healthy range above 30 ng/mL.	Sun exposure, fatty fish
Omega-3 fatty acids (EPA/DHA)	May improve muscle responsiveness to protein and training, partly by supporting normal inflammatory balance and cellular health.	Aim for 2 to 3 servings of omega-3-rich fish per week.	Salmon, sardines, mackerel, cod liver
Magnesium	Required for energy production, vitamin D metabolism, cellular signaling, and normal muscle relaxation.	300 to 400 mg daily.	Nuts, seeds, leafy greens, dark chocolate



Part 3 – Training

This is the step that helps pull everything together: physically training muscles to utilize protein more effectively while simultaneously supporting structural integrity and resilience.

High-intensity resistance and impact training can significantly support strength, bone health, and body composition, and when performed correctly, can be remarkably safe even for older women.

What happened in the study

In a [research trial](#) conducted by *The Journal of Bone and Mineral Research*, postmenopausal women with low bone density, averaging 65 years of age, performed supervised high-intensity resistance and impact training (HiRIT) twice weekly for roughly 30 minutes per session over 8 months.

The protocol required participants to lift heavy weights—performing five sets of five repetitions at 80 to 85 percent of their maximum capacity—alongside brief, impact-based jumping exercises.

Compared to the home-based, low-intensity group, the HiRIT participants showed improvements in several functional strength markers—including bone density and back and leg strength—whereas the low-intensity group actually lost bone mass over the same period.

Furthermore, the study maintained an exceptional safety record, reporting only one minor injury during the trial.

The takeaway: *Lifting heavy and doing impact training—when supervised and progressive—can be both highly supportive and safe for older, thinning bones.*





Important safety note

This is general educational information based on published research, not personal medical advice. Individual needs and results may vary.

Talk to your doctor before starting: If you have osteoporosis, heart problems, high blood pressure, joint issues, or any chronic condition, it is especially important to speak to a professional. Heavy training is powerful, and it should be cleared and, ideally, coached by a certified healthcare or fitness professional.

Start light: The women in the research advanced to heavy weights over months of regular training under close supervision, not on day one. Starting too heavy, too soon can cause injuries and setbacks that undermine the strength and function you are trying to build.

Build up gradually: Always start at a weight that you can control with clean form, and only progress in small increments when your top set feels manageable. Always allow for adequate recovery time between workouts before challenging the same muscle group again.

Pay attention to your body: Feelings of exertion during a challenging set and mild soreness in the day or two after training are normal signs of effort. Sharp pain, joint pain, or lingering pain is not. Stop and seek help for chest pain, dizziness, sharp joint pain, or anything that feels wrong.

It's recommended to consult a professional trainer or physiotherapist to help guide proper form, which is essential for safe and effective training, especially when performing new workout routines.

How to get started:

Follow this phased template to safely build structural tolerance and integrity over time.

Phase 1: Movement mastery (weeks 1–4)

Goal: Master clean form, establish a baseline habit, and wake up dormant muscle fibers.

- Use bodyweight or very light weights

This copy is for informational purposes only and is not intended to diagnose, treat, cure, or prevent any disease. Consult your physician before starting any new health regimen or supplement.



- Aim for 2 to 3 sets of 8 to 10 comfortable reps with each exercise
- Zero-impact landing mechanics practice
- Two sessions a week

Safety note: Impact work can help support strong bones, but if you have osteoporosis, do it only under professional supervision.

Phase 2: Add moderate load (weeks 5–8)

Goal: Condition your connective tissues and bones to handle external resistance.

- Once your form is solid, start adding weight
- Aim for three sets of eight reps—the last 1 to 2 reps of each set feel genuinely challenging, but doable with good form
- Introduce low-impact dynamics, 4 to 5 sets of five reps
- Two sessions a week

Phase 3: Add heavier loads (week 9 and beyond)

Goal: Transition into the actual high-intensity loading parameters. Your form is solid. Weight is comfortably challenging.

- Maintaining clean form, gradually move toward heavier weights at around 80 percent of maximum capacity
- Aim for a total of 4 to 5 sets of five reps—heavier means fewer reps, more rest between sets. NEVER sacrifice your form for a higher weight.
- Introduce impact loading, 4 to 5 sets of five reps
- Two sessions a week



The workout framework

Build every session around one of these resistance exercises—alternating between sessions—followed by the appropriate phase of impact training.

1. Resistance training

Heavy resistance training helps stimulate your muscles to pull harder against your bones, which signals your body to lay down denser, stronger skeletal tissue. It's been associated with improved muscle mass, greater bone mineral density, and a lower risk of frailty in older adults.

Here are the three foundational movements that were used in the trial:

- **Back squats:** This multi-joint movement strengthens the thighs, hips, and lower spine.
- **Deadlifts:** A foundational movement for the back-of-body and everyday strength.
- **Overhead presses:** Builds shoulders, arms, and posture.

2. Impact training finishers

Impact training helps teach your muscles, bones, joints, and tendons how to absorb shock. It's been associated with improved tissue integrity and coordination.

Here is the three-stage progression designed to safely introduce shock absorption to your routine:

- **Zero-impact landing mechanics:** Helps get the body familiar with the movement without your feet ever leaving the floor.
- **Low-impact dynamics:** A gentle way to introduce real loading forces to the skeleton, creating a safe physical vibration that helps stimulate bone-remodeling cells.
- **Impact loading:** The full stimulus utilizes supervised jumping and drop-landings to create the forces needed to target hip bone density



The order of the session

To help maximize your power output and minimize fatigue-driven injury, always perform your workouts in this sequence:

- 1. Warm-Up:** Spend 5 to 10 minutes walking lightly or stretching, followed by 1 to 2 very light-weight practice sets of your lifting movements to lubricate the joints and increase blood flow.
- 2. Resistance training:** Rotate exercises each session
- 3. Impact finish:** Practice different impact movements each session

Quick-start checklist

- ☐ Get your doctor's okay and a vitamin D test before you start
- ☐ Build the daily habit of hitting your protein target within the proper intervals
- ☐ Incorporate supportive nutrients and anchor each meal with a high-leucine food
- ☐ Avoid snacking
- ☐ If fasting, do it sparingly and adjust based on personal needs
- ☐ Start Phase 1 and progress comfortably and gradually over the coming weeks

Continue Your Health Journey, Wherever You Are

Explore our free apps and website for wellness tools at your fingertips

4.8
★ 2.57K


Dr. Berg Junk Food Meter

Make informed food choices
with the tap of a button

The Dr. Berg Junk Food Meter app helps you spot harmful ingredients, ultra-processed additives, and hidden sugars—right from your phone.

Download on the
App Store

GET IT ON
Google Play



5.0
★ 2.46K

The Official Dr. Berg App

Expert health insights,
always within reach

Your go-to resource for educational content on health, wellness, and nutrition. Access a rich library of videos, audio content, and downloadable PDF resources designed to support your personal wellness journey—anytime, anywhere.

Download on the
App Store

GET IT ON
Google Play



Explore more topics to support your health journey at drberg.com



These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease. These nutritional recommendations are supplied for you to share with your health care provider. The content herein and the nutrients recommended are in aid of good health, not disease treatment. Please consult your health care practitioner for diagnosis and treatment of disease.